
Partition Behavior of Perfluorooctane Sulfonate (PFOS) and Perfluorooctanoic acid (PFOA) in Riverine Sediments

R.X. Liu^{*1}, J.Y. Tian^{1,2}, B.Li¹, X.X.Gong² and Y.Y.Liu²

¹ State Key Laboratory of Environmental Criteria and Risk Assessment (SKLECRA), Chinese Research Academy of Environmental Sciences, Beijing 100012, China (Email: liurx@craes.org.cn)

² College of Chemical Engineering, Shandong University of Technology, Zibo 255049, China

Abstract

The sorption of perfluoroalkyl substances (PFASs) on the sediments is an important process which contributes to their fate, distribution and transport in water system. In this study, the sorption kinetic, thermodynamics and effect parameters on partition behavior of perfluorooctane sulfonate (PFOS) and perfluorooctanoic acid (PFOA) at low initial concentration were investigated. The results showed that the equilibrium time was achieved within 48h. The sorption isotherms of PFOS and PFOA could be described by Freundlich equation. The surface area and composition of sediment particles as well as solution pH values and ion strength strongly influenced the sorption of PFOA and PFOS. Both PFOA and PFOS exhibited higher sorption capacity on the sediments with high organic matter content. These results indicated that multiple driving forces such as hydrophobic exclusion, specific chemical interaction and electrostatic attraction contributed to the nonlinear sorption of PFOA and PFOS on sediments.

Keywords

PFOA; PFOS; sediment; partition properties; natural riverine water

INTRODUCTION

Perfluoroalkyl substances (PFASs) are a new type of persistent organic pollutants (POPs) (Wang & Shih 2011; Wang et al. 2012). They have attracted global concern due to high bioaccumulation, extreme persistence and toxicity, as well as wide distribution in the environment. Of the PFASs, perfluorooctane sulfonic acid (PFOS) and perfluorooctanoic acid (PFOA), ubiquitously present in the diverse environments (Beskoski et al. 2013; Du et al. 2014), showed the highest persistence and accumulation in the aquatic environment. Thus, PFOS, its salts and perfluorooctane sulfonyl fluoride (PFOS-F) have been added to the Stockholm Convention list of the POPs since 2009 (Wang et al. 2009). PFOA, as a Substance of Very High Concern (SVHC) in candidate list under REACH regulation (ECHA 2015) has been prohibited by the Environmental Protection Agency of the United State (US-EPA) on a PFOA-stewardship program in 2006(US-EPA 2015).

However, owing to more than 50 years of production and worldwide application, PFOS and PFOA can be released into the natural aquatic environment, causing the global distribution of these compounds (Zhang et al. 2012). Subsequently, these pollutants have undoubtedly involved the complex processes such as sorption, degradation and transport, which determine their environmental fate in aquatic systems (Li et al. 2012). The understanding of these basic mechanisms is of significance in both regulatory and scientific areas. Sediment, with complex compositions including minerals, clay, sand and sedimentary organic matter, is an important sink and reservoir of PFOS and PFOA (Zhao et al. 2014). The sorption of PFOS and PFOA onto the sediment determines their distribution, transport and transformation processes in the aquatic environment. The sorption behavior highly depends on the physical and chemical properties of PFOS and PFOA, as well as the conditions of water chemistry such as pH value, ionic strength (IS), and property of dissolved organic matter (DOM). Higgins and Luthy (2006) evaluated various sediment-, solution-, and chemical-specific parameters potentially affecting sorption of PFASs.

They found that organic carbon rather than inorganic oxide content in sediment was the dominant effect parameter on sorption, indicating the importance of hydrophobic interactions. The similar results were also obtained in other previous studies (Ahrens et al. 2009; Ahrens et al. 2010; Zhao et al. 2012; Zhao et al. 2014; Milinovic et al. 2015), where they reported that hydrophobic interaction predominated the sorption behavior of longer-chain PFASs that were found to adsorb more strongly onto sediment, soil or activated sludge while short-chain PFASs were exclusively in water. In addition, previous work had also demonstrated that sorption of PFASs on sediment, soil or sludge increased with increasing solution Ca^{2+} concentration and decreasing pH value, in particular, the sorption behaviors of PFOS was expected to be strongly affected by solution conditions (Higgins 2006; Zhao et al. 2014; Milinovic et al. 2015). These observations suggested that electrostatic interactions were not negligible in the sorption of PFASs onto sediment. More evidences for both hydrophobic and electrostatic interactions involved in the sorption mechanism of PFASs were found by other researchers using aluminum-rich mineral-humic acid as a simulated sediment and sewage sludges as adsorbents (Zhang et al. 2013; Wang et al. 2015). However, to reduce the detection errors most of these studies were performed with high initial concentrations of PFASs ranging from 1 to 1000 $\mu\text{g/L}$ that were much higher than their concentrations (ng/L) in the natural aquatic environment (Lein et al. 2008; Dufkov áet al. 2012; Zhang et al. 2012; Anumol et al. 2013).

The sorptive behavior of PFASs onto natural aquatic sediment highly depends on many factors. Particularly with high concentration of PFASs to simulate the natural sediment–water interfacial processes is inadequate and possibly leads to absurd conclusions (Ahrens et al. 2011). For this reason, this study examined the sorption behavior of PFOS and PFOA on riverine sediment using a low initial concentration so as to simulate natural aquatic interfacial process. The specific objectives of this study were: by batch partitioning experiments, to determine the sorption coefficient and isotherms of PFOS and PFOA onto the sediment and to examine the effect of pH, ionic strengths, and particle size and density of sediment on the sorption. These would provide supplementary data for understanding natural sediment-water interfacial process of pollutants.

MATERIALS AND METHODS

Reagents and chemicals

PFOA and PFOS were purchased from AccuStandard Inc (USA). $^{13}\text{C}_4$ -labelled PFOA and PFOS (Wellington's laboratory) were used as internal standards. Calcium chloride, sodium azide, hydrochloric acid and sodium hydroxide with analytical reagent grade were purchased from local chemical companies in Beijing, China. HPLC-grade methanol and ammonium acetate were supplied by Fisher Company (USA). Milli-Q ultrapure water was used throughout the experiment. All standard solutions were prepared in methanol and stored in polypropylene (PP) bottles at 4°C.

Sediment collection and characteristics

Four sediment samples were collected from Daliao River systems, Northeast China. The sampling sites were chosen to reflect a variety of physicochemical properties potentially influencing partitioning of PFOA and PFOS. The surface sediments were collected in PP plastic bags using a grab sampler and transferred to the laboratory. The wet samples were air-dried, ground and sieved through 2 mm meshes. Total organic carbon (TOC) in sediments were measured using a high temperature TOC analyser (Dohrmann DC-190) after the removal of inorganic carbon by adding diluted HCl until acidification reaction was completed. Specific surface area (SSA) was analyzed by N_2 sorption method. The pH value of sediment was measured in a 1:2.5 (w/w) mixture of the sediment with 0.01mol/L of CaCl_2 solution by a pH meter. Total Fe and Mn contents in sediment were determined by an ICP-OES Thermo Elemental (TJA) Iris Intrepid spectrometer following digestion of sediment with diluted aqua regia (Li et al. 2012). Sediment particle size fractionation

was conducted by wet sieving the sediments through 40, 80, 120, 240 meshes to obtain three size fractions, i.e. 0.42 ~ 0.20, 0.20 ~ 0.125, 0.125 ~ 0.061 mm. The physicochemical parameters of sediments were shown in Table 1.

Table 1 Physicochemical property of the sediments

Parameters	sediment 1	sediment 2	sediment 3	sediment 4
pH value	7.58	7.38	7.93	7.49
OC (mg/g)	8.09	7.98	7.97	80.4
SSA (m ² /g)	104	66.2	52.1	48.1
Fe (mg/g)	45.8	26.9	19.7	24.3
Mn (mg/g)	0.82	0.38	0.40	0.34

Batch partitioning experiments

Batch experiments included partitioning kinetics, effect of pH value, ion strength and particle size fractions, as well as sorption isotherm experiments. The partitioning experiments were conducted in the 50ml polypropylene copolymer (PPCO) Nalgene centrifuge tubes. Duplicate sets of tubes containing 1.0 g (dry weight) of sediment and 30 ml of 0.01mol/L CaCl₂ and 200 mg/L NaN₃ solution were spiked respectively with a certain amount of PFOA or PFOS, and equilibrated on the Thermostatic shaker at 150 rpm and 25°C. To determine an appropriate equilibration time, the centrifuge tubes with initial PFOA or PFOS concentration of 50 ng/L were shaken for 0 ~ 72 hours and periodically removed at selected time intervals of 0, 0.5, 1, 2, 4, 8, 12, 24, 36, 48, 60 and 72 hours. The effects of pH, ion strength and sediment particle size on partition behavior were investigated by batch sorption equilibration experiment at initial PFOS and PFOA concentrations of 50 ng/L. The pH values of aqueous solution were adjusted in the range of 3 ~ 9 with dilute NaOH or HCl. The concentrations of CaCl₂ were 0.001, 0.01 and 0.1mol/L. The sorption isotherm experiments were carried out with PFOS or PFOA concentrations ranging from 50 to 500 ng/L and at equilibration time of 48 hours. After shaking, the tubes were centrifuged at 1600r/min for 15min and the supernatants were transferred to the new PP tubes for the extraction and analysis.

For each sample batch, blank samples were prepared using 30 mL of millipore water and treated in exactly the same way as the samples. The control experiment in the absence of sediment showed that the total PFOS or PFOA losses was below 5%, which was neglected in the experiment.

Extraction and determination of PFOA and PFOS

The supernatant was passed through Oasis HLB cartridge (Waters, 500mg, 6mL) for the extraction of PFOA and PFOS. The cartridges were firstly cleaned by dichloromethane and methanol respectively to remove residual impurities, subsequently preconditioned by 5 mL each of methanol and ultrapure water (Dufkov á et al. 2012; Gong et al. 2016). After extraction, the cartridges were rinsed with ultrapure water, evacuated to dry for 30 min and eluted with 3×5ml methanol into 15ml PP tubes. The elution gathered was dried by Nitrogen blowing, and then 1mL of methanol-H₂O (v:v, 1:3) and 10µL of 150 µg/L internal standard solution were added for UPLC-MS-MS analysis.

The concentrations of PFOS or PFOA were determined by UPLC-Xevo TQD (Waters, USA). An Acquity UPLC® BEH C₁₈ column (2.1×50 mm, 1.7µm) was used for the separation of analyte. Tandem mass spectrometry was conducted on triple quadrupole mass spectrometer (Xevo TQD, Waters, USA) equipped with an ESI source. The mobile phases were 2 mmol/L of NH₄OAC in water (A) and in methanol (B). The detailed analytical parameters of the target compounds were described in the previous study (Gong et al. 2016).

Sorption data fitting and partition coefficient

Sorption isotherms were constructed by plotting C_{sed} vs. C_w for PFOA or PFOS-sediment batch sorption system. The Freundlich model was used to fit sorption data, as described by equations (1) and (2):

$$C_{sed} = K_f (C_w)^{1/n} \quad (1)$$

or

$$\log C_{sed} = 1/n \log C_w + \log K_f \quad (2)$$

where C_{sed} and C_w are concentrations of PFOA or PFOS in sediment (ng/g) and the solution (ng/L) at equilibrium time; K_f is a Freundlich constant representing the sorption capacity and n is the Freundlich exponent depicting the nonlinearity of sorption.

The interaction of PFOA or PFOS between sediment and water can be described by its partition coefficient (K_d):

$$K_d = C_{sed}/C_w \quad (3)$$

By combination of equations (1) and (3), the K_d values were calculated using the equation (4), which were equilibrium concentration dependent:

$$K_d = K_f (C_w)^{(1-n)/n} \quad (4)$$

Previous studies have shown that the fraction of organic carbon (f_{oc}) has a significant effect on the partition of PFOA or PFOS onto sediment (Higgins and Luthy 2006). Thus, the organic carbon normalised partition coefficient (K_{oc}) was calculated by the following equation (5):

$$K_{oc} = K_d/f_{oc} \quad (5)$$

RESULTS AND DISCUSSION

Sorption kinetics

Figure 1 showed the sorption kinetics of PFOA and PFOS onto sediment. It was obvious that the sorption process could be divided into three apparent stages: a rapid sorption process, a slow sorption process and sorption equilibrium process. The rapid sorption occurred at 0 ~ 10 hours for PFOA and at 0 ~ 5 hours for PFOS. Between 10 (or 5) and 40 hours, the sorption rate for PFOA or PFOS gradually reduced, and then equilibrium was achieved after approximately 40 hours for both chemicals.




Figure 1. Sorption dynamic curve of PFOA and PFOS onto the sediment (25°C)

This result was slightly different with the previous study where equilibrium time for PFOS and PFOA was shorter e.g. 4 hours and 8 hours, respectively, when using higher initial concentration (5 µg/ L) of PFOS and PFOA (Arvaniti et al. 2014). This indicated that the initial concentration of pollutants was one of impact factors controlling their sorptive kinetics between water and sediment. Simulating natural sediment–water interfacial process of pollutants with higher concentration than environmental level would lead to unrealistic results. Therefore, at low concentration of PFOS and PFOA (e.g. 50ng/L) the equilibrium time of 48 hours was adopted for the batch sorption system.

Effect of solution pH

The effect of pH value on the sorption of PFOA and PFOS was shown in Figure 2. It was observed that solution pH values in studied range influenced sorption of PFOA and PFOS onto the sediment to some extent. It was well known that the specific surface species could be formed on the natural sediment surface. Solution pH value would affect specific pH-dependent reactions on sediment surface such as ligand exchange, electrostatic interactions and hydrophobic effect (Zhou et al. 2010; Zhang et al. 2012). Obviously, with increasing pH values between 3 and 7, the sorption of PFOA and PFOS onto the sediment slightly reduced, which could be explained by the electrostatic interaction of PFOA and PFOS anionic species with the positively charged sediment surface at lower pH value. The results were in agreement with Higgins and Luthy’s report (Higgins and Luthy 2006). It was also noted that a slight increase in sorption of PFOS and PFOA on the sediment was found at solution pH over 7. Although more negative charges on sediment surface occurred with pH value increase, resulting in weak electrostatic attraction towards PFOA and PFOS anionic specie, ligand exchange might be predominant on interfacial reactions, leading to enhanced sorption of PFOS and PFOA on the sediment (Zhao et al. 2014).




Figure 2. Effect of pH value on the sorption of PFOS and PFOA

Effect of ion strength

Figure 3 revealed the sorption trends of PFOS and PFOA on the sediment with varying in Ca^{2+} concentration from 0.001 to 0.1 mol/L at solution pH value 7. It indicated that the sorption of PFOS and PFOA increased with increasing Ca^{2+} concentration, which was in accord with previous reports by Higgins and Luthy (2006) and by Chen et al (2009). The effect of Ca^{2+} concentration on increased sorption percentage of PFOS and PFOA was likely due to several reasons. One was linked to reduced solubility of PFOS and PFOA as the ionic strength enhanced, thereby increasing hydrophobic interaction. Chen et al has reported that the sorption-enhancing impact of Ca^{2+}

concentration was possibly related to the concentration of PFOS and PFOA in the aqueous phase with much stronger impact at low PFOS and PFOA concentration (Chen et al. 2009). Furthermore, with the divalent cation Ca^{2+} increasing, the electrostatic repulsion between anionic PFOS or PFOA molecules and the negatively charged sediment surface was reduced because of neutralization of the negative charges on the sediment surface by the high Ca^{2+} concentration, which promoted the sorption of PFOS and PFOA on the sediment. Beyond those Ca-bridging effect initiated by cations in the solution needed to be taken into consideration (Zhao et al. 2014). It was widely reported that divalent cations have been found to be able to shift the negative sites of adsorbent surfaces into positive ones acting as the bridge to electrostatically attract PFASs (Du et al. 2014). You et al. (2010) also observed that sorption of PFOS on the sediment increased when the CaCl_2 concentration increased from 0.005 to 0.5 mol/L, suggesting that Ca-bridging effect was responsible for this result. Therefore, the formation of Ca-bridging with sediment might influence the sorption of PFOA and PFOS on the sediment.




Figure 3. Effect of Ca^{2+} on the sorption of PFOA and PFOS

Effect of particle size

The particle size-dependence of sorption for PFOA and PFOS was shown in Figure 4. The sorption percentage of target contaminants measured at pH 7 decreased with increasing particle size of sediment. The results might be attributed to specific surface areas and physicochemical properties of different particle size. The specific surface areas were decreased with the increased grain size of sediment while the smaller particles would have larger external surface areas (Yu et al. 2009). The diverse compositions and properties of the fractionated sediment, such as organic carbon fraction, debris of wood and stalks, mineral grains, might result in different sorption capacity to PFASs (Zhao et al. 2012).

On the other hand, the compact organic matter was possibly involved with the smallest sediment particles. It has been reported that the organic matter enhanced the sorption capacity of organic pollutants onto sediments (Chen et al. 2009; Beskoski et al. 2013; Li et al. 2014). As shown in Figure 5, the sorption of PFOS and PFOA on sediments was somewhat related to organic matter, indicating that hydrophobic partitioning played a role in the interaction between target pollutants and sediment. Therefore, the combination between small-sized sediment and organic carbon was contributed to the sorption of PFOA and PFOS on sediment, which was consisted with the results reported by Zhao et al. (2012).




Figure 4. Effect of particle size on the sorption of PFOA and PFOS




Figure 5. Effect of organic matter content on sorption of PFOA and PFOS

Sorption isotherms

The sorption isotherms of PFOA and PFOS on sediments were shown in Figure 6. Equilibrium sorption data was calculated based on the Freundlich model. Over the studied concentrations range of PFOA and PFOS, all experimental data was fitted very well by the nonlinear Freundlich type sorption isotherm with the regression coefficients (R^2) of 0.97 for PFOA and 0.95 for PFOS, respectively. For both PFOA and PFOS, the sorption coefficients (K_f) were 0.0051 and 0.0012, and the nonlinearity of sorption ($1/n$) were 0.84 and 0.93, respectively. Regarding nonlinear sorption isotherms, the partition coefficients K_d and K_{oc} values were function of the solute concentration, ranged from 0.0012~ 0.0027 and 0.15~0.36 for PFOA, and 0.00061~ 0.00083 and 0.081~ 0.114 for PFOS.

This nonlinearity of sorption behavior was mainly attributed to organic and mineral matrices involved in sediment. Given that hydrophobicity of perfluorinated chain and hydrophilicity of sulfonate or carboxylate (Zhang et al. 2013), as well as diverse sorption sites on sediments, single driving force seems to be difficult to explain interfacial reaction between pollutants and sediments. The hydrophobic exclusion of perfluorinated chain, specific chemical interaction of sulfonate or carboxylate with sediment, and electrostatic attraction of charged solute molecule with the charged sediment surface might contribute to the nonlinear sorption of PFOA and PFOS, forming the sorption of multi-molecular layers.




Figure 6. Sorption isotherms of PFOS and PFOA on the sediments

CONCLUSIONS

In natural aquatic system, the sorption of PFOS and PFOA onto the sediments was very low. The kinetic experiments showed that the equilibrium time was achieved within 48h. The sorption behavior of PFOA and PFOS could be described by Freundlich equation. Both PFOA and PFOS displayed higher sorption capacity onto the sediments at high ionic strength and pH values. The sediment particle with small size and high organic matter content enhanced the sorption of PFOA and PFOS due to the surface area and specific composition of particles. These results had important environmental implications: the leaching of PFCs from sediment would become more serious if background solution was neutral and with low ionic strength. The research would be benefit for the efficient removal of PFOS and PFOA in the natural aquatic systems.

ACKNOWLEDGEMENTS

This work was supported by the National Natural Science Foundation of China (No.21277133) and the Basic Scientific and Research Program for Central Nonprofit Research Institutes (No.2012-YSKY-11)

REFERENCE

Ahrens, L., Taniyasu, S., Yeung, L. W., Yamashita, N., Lam, P. K., Ebinghaus, R. 2010. Distribution of polyfluoroalkyl compounds in water, suspended particulate matter and sediment from Tokyo Bay, Japan. *Chemosphere* **79**(3), 266-272.

-
- Ahrens, L., Yamashita, N., Yeung, L. W., Taniyasu, S., Horii, Y., Lam, P. K., Ebinghaus, R. 2009. Partitioning behavior of per-and polyfluoroalkyl compounds between pore water and sediment in two sediment cores from Tokyo Bay, Japan. *Environmental science & technology* **43**(18), 6969-6975.
- Ahrens, L., Yeung, L. W., Taniyasu, S., Lam, P. K., Yamashita, N. 2011. Partitioning of perfluorooctanoate (PFOA), perfluorooctane sulfonate (PFOS) and perfluorooctane sulfonamide (PFOSA) between water and sediment. *Chemosphere* **85**(5), 731-737.
- Anumol, T., Merel, S., Clarke, B. O., Snyder, S. A. 2013. Ultra high performance liquid chromatography tandem mass spectrometry for rapid analysis of trace organic contaminants in water. *Chemical Central Journal* **7**, 104.
- Arvaniti, O. S., Andersen, H. R., Thomaidis, N. S., Stasinakis, A. S. 2014. Sorption of Perfluorinated Compounds onto different types of sewage sludge and assessment of its importance during wastewater treatment. *Chemosphere* **111**, 405-411.
- Beskoski, V. P., Takemine, S., Nakano, T., Slavkovic Beskoski, L., Gojgic-Cvijovic, G., Ilic, M., Miletic S., Vrvic, M. M. 2013. Perfluorinated compounds in sediment samples from the wastewater canal of Pancevo (Serbia) industrial area. *Chemosphere* **91**(10), 1408-1415.
- Chen, H., Chen, S., Quan, X., Zhao, Y., Zhao, H. 2009. Sorption of perfluorooctane sulfonate (PFOS) on oil and oil-derived black carbon: influence of solution pH and $[Ca^{2+}]$. *Chemosphere* **77**(10), 1406-1411.
- Du, Z., Deng, S., Bei, Y., Huang, Q., Wang, B., Huang, J., Yu, G. 2014. Adsorption behavior and mechanism of perfluorinated compounds on various adsorbents--a review. *Journal Hazardous Materials* **274**, 443-454.
- Dufková V., Čabala, R., Ševčík, V. 2012. Determination of C 5–C 12 perfluoroalkyl carboxylic acids in river water samples in the Czech Republic by GC–MS after SPE preconcentration. *Chemosphere* **87**(5), 463-469.
- ECHA 2015. Candidate List of substances of very high concern for Authorisation. <http://echa.europa.eu/web/guest/candidate-list-table>.
- Gong, X., Liu, R., Li, B., Song, Y., Liu, Y. 2016. Perfluoroalkyl acids in Daliao River system of northeast China: determination, distribution and ecological risk. *Environmental Earth Sciences* **75**(6), 1-10.
- Higgins, C. P., Luthy, R.G. 2006. Sorption of Perfluorinated on sediments. *Environment Science & Technology* (40), 7251-7256.
- Lein, N. P. H., Fujii, S., Tanaka, S., Nozoe, M., Tanaka, H. 2008. Contamination of perfluorooctane sulfonate (PFOS) and perfluorooctanoate (PFOA) in surface water of the Yodo River basin (Japan). *Desalination* **226**(1), 338-347.
- Li, C., Ji, R., Schaffer, A., Sequaris, J. M., Amelung, W., Vereecken, H., Klumpp, E. 2012. Sorption of a branched nonylphenol and perfluorooctanoic acid on Yangtze River sediments and their model components. *Journal Environmental Monitoring* **14**(10), 2653-2658.
- Li, Y., Zhang, C., Li, S., Zhou, C., Li, X. 2014. Single and competitive adsorption of 17alpha-ethinylestradiol and bisphenol A with estrone, beta-estradiol, and estriol onto sediment. *Marine Drugs* **12**(3), 1349-1360.
- Milinic, J., Lacorte, S., Vidal, M., Rigol, A. 2015. Sorption behaviour of perfluoroalkyl substances in soils. *Science of the Total Environment* **511**, 63-71.
- US-EPA Environmental Protection Agency 2015. 2010/2015 PFOA Stewardship Program.

<http://www.epa.gov/oppt/pfoa/pubs/stewardship/index.html>.

- Wang, F., Liu, C., Shih, K. 2012. Adsorption behavior of perfluorooctanesulfonate PFOS and perfluorooctanoate (PFOA) on boehmite. *Chemosphere* **89**(8), 1009-1014.
- Wang, F., Shih, K. 2011. Adsorption of perfluorooctanesulfonate (PFOS) and perfluorooctanoate (PFOA) on alumina: influence of solution pH and cations. *Water Research* **45**(9), 2925-2930.
- Wang, F., Shih, K., Leckie, J. O. 2015. Effect of humic acid on the sorption of perfluorooctane sulfonate (PFOS) and perfluorobutane sulfonate (PFBS) on boehmite. *Chemosphere* **118**, 213-218.
- Wang, T., Wang, Y., Liao, C., Cai, Y., Jiang, G. 2009. Perspectives on the Inclusion of Perfluorooctane Sulfonate into the Stockholm Convention on Persistent Organic Pollutants 1. *Environmental Science and Technology* **43**(14), 5171-5175.
- You, C., Jia, C., Pan, G. 2010. Effect of salinity and sediment characteristics on the sorption and desorption of perfluorooctane sulfonate at sediment-water interface. *Environmental Pollution* **158**(5), 1343-1347.
- Yu, Q., Zhang, R., Deng, S., Huang, J., Yu, G. 2009. Sorption of perfluorooctane sulfonate and perfluorooctanoate on activated carbons and resin: Kinetic and isotherm study. *Water Research* **43**(4), 1150-1158.
- Zhang, C., Yan, H., Li, F., Hu, X., Zhou, Q. 2013. Sorption of short-and long-chain perfluoroalkyl surfactants on sewage sludges. *Journal Hazardous Materials* **260**, 689-699.
- Zhang, Y., Meng, W., Guo, C., Xu, J., Yu, T., Fan, W., Li, L. 2012. Determination and partitioning behavior of perfluoroalkyl carboxylic acids and perfluorooctanesulfonate in water and sediment from Dianchi Lake, China. *Chemosphere* **88**(11), 1292-1299.
- Zhao, L., Bian, J., Zhang, Y., Zhu, L., Liu, Z. 2014. Comparison of the sorption behaviors and mechanisms of perfluorosulfonates and perfluorocarboxylic acids on three kinds of clay minerals. *Chemosphere* **114**, 51-58.
- Zhao, L., Zhu, L., Yang, L., Liu, Z., Zhang, Y. 2012. Distribution and desorption of perfluorinated compounds in fractionated sediments. *Chemosphere* **88**(11), 1390-1397.
- Zhou, Q., Deng, S., Zhang, Q., Fan, Q., Huang, J., Yu, G. 2010. Sorption of perfluorooctane sulfonate and perfluorooctanoate on activated sludge. *Chemosphere* **81**(4), 453-458.