

Influence of mineral-organic fertilizers on soil microbial respiration and nitrifying bacteria activity

R. Jarosz¹, L. Marcińska-Mazur¹, J. Szerement¹, M. Juda¹, J. Mokrzycki¹, K. Gondek², M. Mierzwa-Hersztek^{1,2}

¹Department of Mineralogy, Petrography and Geochemistry, AGH University of Science and Technology, al. Mickiewicza 30, 30-059 Krakow, Poland

²Department of Agricultural and Environmental Chemistry, University of Agriculture in Krakow, Mickiewicza 21, 31-120 Krakow

*corresponding authors e-mails: marcinsk@agh.edu.pl;

Key words: mineral-organic fertilizers, soil microbial respiration, nitrifying bacteria activity

Undoubtedly, the amount of waste produced is increasing year by year. Therefore, it is very important to search for new possibilities of transforming and reusing waste materials. One such type of waste is fly ash, which is generated as a by-product in conventional power generation in coal-fired power plants. As it turns out, fly ash contains significant amounts of valuable components that can be reused. The synthesis process can produce zeolites – which are aluminosilicates with specific properties (Belviso 2018; Czarna-Juszkiewicz *et al.* 2020). Zeolites contain a complex system of channels and chambers in their structure, due to which they possess molecular-sieve properties. The properties of zeolites such as porosity, high ion exchange capacity and selectivity are very important from agricultural point of view (Ramesh and Reddy 2011). The combination of zeolites with minerals and the source of external organic matter in the form of mineral-organic fertilizer may lead to an optimized use of nutrients (Cairo *et al.* 2017; Rosalina *et al.* 2019).

One of the methods of assessing the impact of the use of mineral-organic fertilizers with the addition of this materials is the microbiological activity of the soil. In order to determine the effect of applying innovative mineral-organic fertilizers with the addition of zeolites on soil microbial activity, we conducted a pot experiment in a vegetation hall using spring wheat as a test plant. The scheme of the experiment including: C (control soil without any fertilizers); MF (soil with only mineral fertilizers -NPK); CW3% (soil with addition of 3% of lignite and 3% of zeolite-vermiculite composite - NaX-Ver) ; CW6% (soil with addition of 6% of lignite and 3% zeolite-vermiculite composite - NaX-Ver); CL3% (soil with addition of 3% of leonardite and 3% of zeolite-vermiculite composite - NaX-Ver); CL6% (soil with addition of 6% of leonardite and 3% of zeolite-vermiculite composite - NaX-Ver). Each treatment was performed in 4 replications.


Soil for biochemical analyses was collected 5 months after of the mineral-organic mixtures application and stored at 4 °C for biological analysis and at 25°C for chemical analysis. Soil microbial respiration (BR – basal respiration; SIR – substrate induced respiration; QR – respiratory activation quotient of control and amended soil) and nitrifying bacteria activity analysis was performed to determine the biological activity of the soil.

Picture 1. The course of the analysis Soil Microbial Respiration




The soil respiration is referred to metabolic activity of microorganisms. This parameter depends on the physiological condition of soil microorganisms and other environmental factors (Mierzwa-Hersztek *et al.* 2018). The nitrification potential enables determination of the potential activity of a specialised group of autotrophic bacteria— ammonia-oxidising bacteria responsible for the first phase of nitrification (Mierzwa-Hersztek *et al.* 2018).

Figure 1. Basal respiration (BS) and substrate induced respiration (SIR)




The significantly highest BR value was determined in control soil without fertilization (Fig. 1). While the values obtained in the objects: CW3%, CW6%, CL3% and CL6% were on average 21% lower than in the control treatment. The addition of a substrate (glucose) to the analysed soil samples caused a considerable increase in the respiration rate (Fig. 1). The highest SIR values were obtained in the CL6% treatment.

Figure 2. The respiratory-activation quotient (QR)



The respiratory activation quotient (QR) (Fig. 2) that was calculated in our study represents the number of dormant or active microorganisms. The highest values of QR were determined in the soil without fertilizers (C), while in the other treatments the values of this parameter were significantly lower. The lowest QR value (0.23) were obtained in CL3% and CL6% treatments.

Figure 3. Determination of potential nitrification



Our study shows that the addition of 6% lignite and 3% zeolite-vermiculite composite had significant effect on nitrifying bacteria activity (Fig. 3). Compared to the C treatment, nitrifying bacteria activity was higher 40% on average, and compared to the MF treatment 15% on average.

Conclusion: The nitrifying bacteria activity determined in the studied soils indicated that the most favorable conditions for the course of nitrification occurred in the object CW6%. The use of mineral-

organic fertilizers increased the SIR value.

References:

1. Belviso C. 2018. Ultrasonic vs hydrothermal method: Different approaches to convert fly ash into zeolite. How they affect the stability of synthetic products over time. *Ultrason. Sonochem.*, 43, 9-14.
2. Czarna-Juszkiewicz D., Kunecki P., Panek R., Madej J., Wdowin M. 2020. Impact of fly ash fraction on the zeolitization process, *Materials* 2020, 13, 1035; doi:10.3390/ma13051035.
3. Ramesh K., Reddy D.D. 2011. Zeolites and Their Potential Uses in Agriculture [in:] *Advances in Agronomy*, Volume 113: 219-241, DOI: 10.1016/B978-0-12-386473-4.00004-X.
4. Cairo P., de Armas J.M., Artiles P.T., Martin B.D., Carrazana R.J., Lopez O.R. 2017. Effects of zeolite and organic fertilizers on soil quality and yield of sugarcane. *AJCS* 11(06):733-738 (2017) doi: 10.21475/ajcs.17.11.06.p501.
5. Rosalina, F.; Gafur, M.A.A.; Irnawati, I.; Soekamto, M.H.; Sangadji, Z.; Kahar, S.M. 2019. Utilization of compost and zeolite as ameliorant on quartz sand planting media for caisim (*Brassica juncea*) Plant growth. *J. Phys. Conf. Ser.* 2019, 1155, 012055.
6. Mierzwa-Hersztek M., Klimkowicz-Pawlas A., Gondek K. 2018. Influence of Poultry Litter and Poultry Litter Biochar on Soil Microbial Respiration and Nitrifying Bacteria Activity. *Waste Biomass Valor* (2018) 9:379–389 DOI 10.1007/s12649-017-0013-z.

Acknowledgements: Article was conducted under the project “Fly ashes as the precursors of functionalized materials for applications in environmental engineering, civil engineering and agriculture” - project is carried out within the TEAM-NET programme of the Foundation for Polish Science POIR.04.04.00-00-14E6/18-00